



## Dermosporin<sup>®</sup>-β Cream

(Clotrimazole and Betamethasone Dipropionate Cream USP)

ڈرمواسپورن<sup>®</sup>-بی کریم  
(کلوٹرازیمل اور بیٹا میتھاسون)

### COMPOSITION:

Each gram contains:  
Clotrimazole USP ... 10mg.  
Betamethasone Dipropionate USP eq. to Betamethasone 0.5mg.  
[USP Specs.]

### ACTION:

**Dermosporin-β** Cream combines the broad-spectrum antifungal activity of Clotrimazole with sustained anti-inflammatory, antipruritic and vasoconstrictive actions of Betamethasone dipropionate. Clotrimazole appears to act on the fungal cell membrane, causing leakage of cell contents.

### INDICATIONS:

**Dermosporin-β** Cream is indicated for the topical treatment of the following dermal infections: Tinea pedis. Tinea cruris, Tinea corporis due to Trichophyton rubrum, Trichophyton mentagrophytes, Epidermophyton floccosum and Mycosporum canis; Candidiasis due to Candida albicans.

### CONTRA-INDICATIONS:

**Dermosporin-β** is contraindicated in those patients with a history of sensitivity reactions to any of its ingredients, other corticosteroids or imidazoles.

### DOSAGE & ADMINISTRATION:

Gently massage sufficient **Dermosporin-β** Cream into the affected and surrounding skin areas twice daily, in the morning and evening for two weeks in tinea cruris, tinea corporis and candidiasis and for four weeks in tinea pedis.

### OVER DOSAGE:

Symptoms: excessive or prolonged use of topical corticosteroids can suppress pituitary-adrenal function, resulting in secondary adrenal insufficiency and produce manifestations of hypercorticalism, including Cushing's disease.

**Treatment:** In case of chronic toxicity, slow withdrawal of corticosteroids is advised.

Rev. 02-21/2

**PRECAUTIONS:**

**Dermosporin-β** Cream should not be used with occlusive dressing. If irritation or sensitization develops with the use of **Dermosporin-β** Cream, treatment should be discontinued and appropriate therapy should be instituted. In the presence of bacterial infection an appropriate antibacterial agent should be administered concomitantly. If a favorable response does not occur promptly **Dermosporin-β** cream should be discontinued until the infection has been controlled adequately.

**ADVERSE REACTIONS:**

The following adverse reactions have been reported infrequently with Clotrimazole and Betamethasone dipropionate when used in combination: paresthesia, maculopapular rash, edema and secondary infections. Reported adverse reactions to Clotrimazole include erythema, stinging, blistering, peeling, edema, pruritus, urticaria and general irritation of the skin. The following local adverse reactions have been reported within the use of topical corticosteroids: burning, itching, irritation, dryness, folliculitis, hypertrichosis, acneiform eruptions, hypo-pigmentation, perioral dermatitis, contact dermatitis, maceration of the skin, secondary infections and skin atrophy.

**INSTRUCTIONS:**

**For external use only.** Store below 30°C. Protect from heat & light. Keep out of the reach of children.

**PRESENTATION:**

**Dermosporin-β** Cream is available in pack sizes of 10g & 15g.

ہدایات: صرف بیرونی استعمال کے لئے۔ سو ڈگری سینٹی گریڈ سے کم درجہ حرارت پر رکھیں۔  
گرمی اور روشنی سے بچائیں۔ بچوں کی پہنچ سے دور رکھیں۔



Manufactured by:  
**NABIQASIM INDUSTRIES (PVT.) LTD.**  
17/24, Korangi Industrial Area,  
Karachi-Pakistan.

Rev: 02-21/2