

## **GLANDIN-E2** Vaginal **(DINOPROSTONE Eur. P.)** Tablet

### **COMPOSITION:**

Each Vaginal tablet contains:  
Dinoprostone Eur. P. ... 3mg.  
NQ's Specs.

### **INDICATIONS:**

Oxytocic agent. GLANDIN-E2 Vaginal tablet is indicated for the induction of labour when there are no maternal or foetal contra-indications.

Clinical safety and efficacy has been established for labour induction with following diagnosis:

Elective labour induction.  
Postmaturity.  
Hypertensive disorders.  
Retardation of foetal development.  
Poor obstetric history.  
Mother over 35 years.  
RH disease.  
Diabetes mellitus (latent/manifest).

### **DOSAGE AND ADMINISTRATION:**

One tablet (3mg) to be inserted into the posterior fornix. A second vaginal tablet may be inserted after 6-8 hours if labour is not established. Maximum dose 6mg (2 tablets).

### **CONTRA-INDICATIONS:**

There is no absolute contra-indication to the use of GLANDIN-E2 Vaginal tablet. However it is not recommended in the following conditions:

When the patient is hypersensitive to prostaglandins.

Patient with ruptured foetal membranes.  
Cases with history of pelvic inflammatory disease.

For patients in whom oxytocic drugs are generally contra-indicated or where prolong contraction of uterus are considered inappropriate such as:

Cases with history of caesarean section or major uterine surgery. Cases with major degree of cephalopelvic disproportion. Cases with foetal distress. Cases with history of difficult labour and / or traumatic delivery.

Grand multiparae with 6 or more previous term pregnancies.

### **WARNING AND PRECAUTIONS:**

Since it has been found that prostaglandins may potentiate the effect of oxytocin, it is recommended that, if these drugs are used together or in sequence, the patient's uterine activity should be carefully monitored. During its use uterine activity, foetal status, and the progression of cervical dilation should be carefully monitored to detect possible evidence of undesired responses, e.g. hypertonus, sustained uterine contractions, or foetal distress. In cases where there is a known history of hypertonic uterine contractility or tetanic uterine contractions, it is recommended that uterine activity and the state of the foetus should be continuously monitored throughout labour. The possibility of uterine rupture should be borne in mind where high tone myometrial contractions are sustained. In labour induction foeto-pelvic relationships should be carefully evaluated before the use of GLANDIN-E2 Vaginal tablet.

Rev: 01-18/2

Caution should be exercised in the administration of GLANDIN-E2 Vaginal tablet for the induction of labour in patients with: Asthma or history of asthma. Glaucoma or raised intraocular pressure. There is no evidence that short term administration of prostaglandin E2 can cause similar bone effects.

**SIDE EFFECTS:**

Very occasional nausea and vomiting. Uterine hypertonus or unduely severe uterine contractions have rarely been encountered.

**OVER DOSAGE:**

Treatment of overdosage is symptomatic.

**INSTRUCTIONS:**

Not to be taken orally.  
Store between 2 ~ 8°C.  
Keep out of the reach of children.  
To be used only under the doctor's supervision.

**PRESENTATION:**

GLANDIN-E2 Vaginal tablet is available in pack of 1 Vaginal tablet.

گلیڈن این ای ۲ فرج کی گولی  
(ڈائنو پروسٹون)

اجزائے ترکیب:

ہر گولی میں ۳ ملی گرام ڈائنو پروسٹون موجود ہے۔

فوائد:

یہ دردزہ کو پیدا کرنے کے لئے ہے بشرطیکہ ماں یا بچے میں اس دوا کے خلاف کوئی ممنوع علامات نہ پائی جائیں۔

ہدایات:

دوا کھانے کے لئے نہیں ہے۔

دوا کو ۲ سے ۸ درجہ سینٹی گریڈ کے درمیان رکھیں۔

بچوں کی پہنچ سے دور رکھیں۔

صرف ڈاکٹر کی نگرانی میں استعمال کریں۔

طریقہ فراہمی:

گلیڈن این ای ۲ فرج کی گولی صرف ایک فرج کی گولی کے پیک میں سپلائی کی جاتی ہے۔



PHARMA

Manufactured by:

**NABIQASIM INDUSTRIES (PVT) LTD.**

17/24, Korangi Industrial Area,  
Karachi-Pakistan.

Rev: 01-18/2