

GYNOSPORIN[®] 500mg Vaginal Tablet / 2% Vaginal Cream / -1 Vaginal Cream (Clotrimazole Vaginal Cream / Vaginal Inserts USP)

COMPOSITION:

GYNOSPORIN Vaginal Tablet:

Each vaginal insert contains:
Clotrimazole USP ... 500mg, [USP Specs.]

GYNOSPORIN Vaginal Cream:

Each gram cream contains:
Clotrimazole USP ... 20mg [USP Specs.]

GYNOSPORIN-1 Vaginal Cream:

Each gram cream contains:
Clotrimazole USP ... 100mg [USP Specs.]

DESCRIPTION: Clotrimazole is broad spectrum antifungal agent for fungal infections. It also exhibits activity against Trichomonas, Staphylococci, Streptococci and Bacteroids. It has no effect on Lacto-bacilli.

INDICATIONS: It is indicated in Candida Vaginitis, Vulvitis and mixed vaginal infections. In case of Trichomonas, **GYNOSPORIN** is recommended along with other oral therapy.




CONTRA-INDICATION: Hypersensitivity to Clotrimazole or to any other component of cream / tablet.

DOSAGE & ADMINISTRATION:

GYNOSPORIN Vaginal Tablet: One tablet should be inserted with applicator into the vagina as high as possible preferably before going to sleep at night.

How to use GYNOSPORIN Vaginal Tablet: First of all wash your hands and dry.

- 1) Remove the applicator from the packaging. Pull out the plunger A until it stops. Remove the tablet from the blister foil pack and place firmly into the applicator B in such position that bullet shape of tablet is towards outside.
- 2) To fit the tablet into the applicator, the holder of the applicator needs to be squeezed lightly at both sides.
- 3) Carefully put the applicator as deep as is comfortable into the vagina (this is easiest when lying on your back with your knees bent up). Holding the applicator in place, slowly press the plunger until it stops so that the tablet is deposited into the vagina.
- 4) Remove the applicator and dispose it.






GYNOSPORIN Vaginal Cream 2%: Insert one filled applicator (5g cream) deeply intravaginally twice daily for three consecutive days or once daily for 6 consecutive days preferably before going to sleep at night.

GYNOSPORIN-1 Vaginal Cream: Insert one filled applicator (5g cream) deeply intravaginally (as a single dose therapy), preferably before going to sleep at night.

How to use GYNOSPORIN Vaginal Cream: First of all wash your hands and dry.

- 1) Before use, pierce the tube seal by inverting the cap over the end of the tube and fix the applicator on to the mouth threads of tube.
- 2) Release cream in the applicator by squeezing the tube gently until plunger stops. Remove the applicator from the mouth of tube by unscrewing and replace the cap on the tube.
- 3) Carefully put the applicator as deep as is comfortable into the vagina (this is easiest when lying on your back with your knees bent up). Holding the applicator in place, slowly press the plunger until whole dose of the cream is transferred into the vagina.
- 4) Remove the applicator carefully. Maintain the position for sometime so that the medicine gets absorbed properly. The applicator may be use again after washing and drying.



Candida Vulvitis or Vulvo-Vaginitis: For candida vulvitis the **GYNOSPORIN** Vaginal Cream 2% should be applied thinly to the external genitalia 2 or 3 times a day for 1-2 weeks.

Rev: 02-22/11

کامینو اسپورن (کلورائیڈز آل پوائس نی)

گائے منہ اسوار :- اس کا کریم سے بھرا ہوا انتہائی کم (۵ گرام) صرف ایک دفعہ اندام نہانی میں داخل کر لیں۔ سب سے پہلے ہاتھ دھو کر خشک کر لیں۔

ممنوعات: کلوزٹرائزڈ آل سے زود حساسیت یا کسی دوسرے جوتزہ کیبی سے حساسیت۔ اندام نہانی کے علاج (vaginal use) کے لئے عمل کے پہلے تین مہینوں کی مدت کے دوران خاص طور سے محتاط رہنا چاہئے۔

گائینو اسپورن ۲۲ ویجیٹل کریم ۳۵ گرام ٹیوب میں ایک اعلیٰ کلاس کے ساتھ دستیاب ہے۔

گامینو اسپورن۔ اویجائل کریم ۵ گرام ٹیوب میں ایک اسپیکیکلر کے ساتھ دستیاب ہے۔

Rev: 02-22/11

Manufactured by:
NABIQASIM INDUSTRIES (PVT.) LTD.
17/24, Korangi Industrial Area, Karachi-Pakistan.